

## LPRO-GDGT

### Description:

Modified GDGT from *Sulfolobus acidocaldarius* for investigational use only.

### Sample information:


Product name	L-prolinol-GDGT
Cat.#	87155
Physical appearance	slightly yellow oil
Apparent pKa value	6.67
Solubility in ethanol	20 mg/mL
Solubility in <i>i</i> -propanol	100 mg/mL
Shipment	ambient temperature, packed under N <sub>2</sub>
Storage	-20 °C

### Sample composition:

Lipid component	Chemical formula	Purity <sup>2</sup>	Molecular mass (g/mol)
LPRO-GDGT <sup>1</sup>	C <sub>96</sub> H <sub>182</sub> N <sub>2</sub> O <sub>6</sub>	>95%	1460.52

GDGT... glycerol dialkyl glycerol tetraether

### Structure:



### Handling information:

Recommended solvents: dissolves in all common organic solvents (e.g. diethylether, dichloromethane, chloroform, THF, *i*-propanol, DMSO...)

The compound is stored under N<sub>2</sub> atmosphere.

For formulation experiments ethanol *absolute* or pure *i*-propanol are recommended as solvent. Note that traces of water, e.g. due to usage of ethanol 96%, lead to formation of a cloudy suspension.

To quantitatively dissolve the product in the original container it is recommended to thoroughly rinse the whole vial and cap with solvent.

<sup>1</sup> The GDGT moiety naturally occurs with 0 to 8 cyclopentane rings, resulting in minor deviations of the molecular mass.

<sup>2</sup> Based on NMR